



ANOGEN - A Division of YES Biotech Laboratories Ltd.

2355 Derry Road East, Unit 23, Mississauga, ON, Canada L5S 1V6 • Tel: (905) 677-9221 • Fax: (905) 677-0023

Mouse Anti-Human MCP-1 Monoclonal Antibody Datasheet

Product Name: mAb anti-Human MCP-1

Clone No.: 9G10

Catalogue No.: MO-C40021E

Quantity: 0.5 mg/vial

Description: Mouse monoclonal antibody to human Monocyte Chemotactic Protein-1 (MCP-1)/ Monocyte Chemotactic and Activating Factor (MCAF)

Purification: Protein G affinity purified

Target Protein: Human MCP-1

Immunogen: Purified recombinant human MCP-1

Fusion Sp2/O-Ag14

Myeloma:

Specificity: This antibody reacts with natural and recombinant human MCP-1.

Species Human, others not tested

Reactivity:

Cross-reactivity: This antibody does not react with human interleukin-8 (IL-8) and other human cytokines tested such as interleukin-1 β (IL-1 β), serum amyloid A (SAA) and epidermal growth factor (EGF).

Host / Isotype: Mouse, IgG1 Kappa

Formulation: Lyophilized from a solution in 0.01M PBS, pH 7.2

Reconstitution: Double distilled water is recommended to adjust the final concentration to 1.00mg/mL.

Storage: Store at -20°C.

Research Area: Immunology, cytokine and chemotaxis

Background: Monocyte chemotactic and activating factor (MCAF) is also called monocyte chemotactic protein-1 (MCP-1) and chemokine (C-C motif) ligand 2 (CCL2). It is primarily secreted by monocytes, macrophages and dendritic cells. This cytokine displays chemotactic activity for monocytes, T-cells, and basophils, but not for neutrophils or eosinophils. MCAF causes the degranulation of basophils and mast cells, and augments the activity of monocyte and macrophage. MCAF plays an important role in inflammation, angiogenesis, autoimmune diseases, renal diseases, chronic infection and granuloma formation.

Applications: **ELISA:** React with human MCP-1.

Western Blot: 0.1 μ g/lane of recombinant human MCP-1 can be detected when antibody is used at 0.1-1 μ g/mL.

Neutralizing: The antibody neutralized 1nM recombinant human MCP-1 induced monocyte chemotaxis in blind well chemotaxis chambers.

References: If research is published using this product, please inform Anogen in order to cite the reference on this datasheet. Anogen will provide one unit of product in the same category as gratitude.

This product is for **LABORATORY RESEARCH USE** and further manufacture **ONLY**, and cannot be administered to human and animals for use in diagnostic and therapeutic procedures.

Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.