

**Mouse Anti-Human SAA Monoclonal Antibody Datasheet****Product Name:** mAb anti-Human SAA**Clone No.:** 607**Catalogue No.:** MO-C40028E**Quantity:** 0.5 mg /vial**Description:** Mouse monoclonal antibody to human serum amyloid A (SAA)**Purification:** Protein G affinity purified**Product type:** Primary antibody - **Detection** antibody in matched antibody pair.**Target Protein:** Human SAA**Immunogen:** Highly purified recombinant human SAA (MW: 12 kDa)**Fusion** Sp2/0-Ag14**Myeloma:****Specificity:** This monoclonal antibody reacts with natural and recombinant human SAA.**Species** Human, others not tested**Reactivity:****Cross-Reactivity:** Does not show any cross-reaction with other human cytokines or growth factors tested such as IL-1 β , IL-8, MCAF, TGF- β and EGF.**Host / Isotype:** Mouse, IgG2a Kappa**Formulation:** Lyophilized from 0.01M K₂HPO₄ – KH₂PO₄ buffer, pH = 7.0**Reconstitution:** Double distilled water is recommended to reconstitute the antibody.**Storage:** Store at -20°C**Research Area:** Acute phase protein, Inflammatory reaction**Background:** SAA is an acute phase protein and a sensitive inflammatory marker. Cytokines such as IL-1, IL-6 and TNF- α stimulates hepatocytes to produce and release SAA. In acute phase, SAA in blood can increase to more than 1,000 folds of normal level. Prolonged and repeated elevation of SAA results in abnormal deposition of amyloid proteins in organ or tissue in insoluble beta pleated form. Determination of SAA levels is important for assessment of renal allograft rejection, estimation of tissue damage caused by myocardial infarction and severe angina, and diagnosis of inflammation in cystic fibrosis patients and in patients infected with virus.**Applications:** **ELISA:** This antibody can be used as a detection antibody in sandwich ELISA applications for human SAA detection in combination with a capture antibody (Clone 115, Cat. No.: C40028A)**Western Blot:** The antibody, when used at concentration of 0.1-0.5 μ g/mL will allow visualization of 100ng/lane of recombinant human SAA.

This product is for **LABORATORY RESEARCH USE** and further manufacture **ONLY**, and cannot be administrated to human and animals for use in diagnostic and therapeutic procedures.

Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.



ANOGEN - A Division of YES Biotech Laboratories Ltd.

2355 Derry Road East, Unit 23, Mississauga, ON, Canada L5S 1V6 • Tel: (905) 677-9221 • Fax: (905) 677-0023

References:

1. Wu TL et al. Establishment of an in-house ELISA and the reference range for serum amyloid A (SAA): complementarity between SAA and C-reactive protein as markers of inflammation. 2007 Feb; Clin Chim Acta. 376(1-2):72-6.

2. Chang Yeop Han et al. Differential Effect of Saturated and Unsaturated Free Fatty Acids on the Generation of Monocyte Adhesion and Chemotactic Factors by Adipocytes. Diabetes. Feb 2010; 59(2): 386-396.

If research is published using this product, please inform Anogen in order to cite the reference on this datasheet. Anogen will provide one unit of product in the same category as gratitude.

This product is for LABORATORY RESEARCH USE and further manufacture ONLY, and cannot be administered to human and animals for use in diagnostic and therapeutic procedures.

Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.